

NOAA Technical Memorandum NWS WR- 143

THE DEPTH OF THE MARINE LAYER AT SAN DIEGO AS RELATED TO  
SUBSEQUENT COOL SEASON PRECIPITATION EPISODES IN ARIZONA

Ira S. Brenner  
National Weather Service Forecast Office  
Phoenix\*, Arizona

May 1979

\*Present affiliation: National Hurricane Center  
Miami, Florida

UNITED STATES  
DEPARTMENT OF COMMERCE  
Juanita M. Kreps,  
Secretary

NATIONAL OCEANIC AND  
ATMOSPHERIC ADMINISTRATION  
Richard Frank,  
Administrator

NATIONAL WEATHER  
SERVICE  
Richard E. Hallgren,  
Director



This Technical Memorandum has been reviewed and is approved for publication by Scientific Services Division, Western Region.

A handwritten signature in black ink, appearing to read "L. W. Snellman". The signature is written in a cursive style with a long, sweeping tail that extends to the right.

L. W. Snellman, Chief  
Scientific Services Division  
Western Region Headquarters  
Salt Lake City, Utah

## CONTENTS

	<u>Page</u>
Figures . . . . .	iv
Abstract . . . . .	1
I. Introduction . . . . .	1
II. Developmental Data Base Stratification . . . . .	2
III. Developmental Data Sample Analysis . . . . .	3
IV. Independent Test Data . . . . .	5
V. Conclusions . . . . .	7
VI. Acknowledgments . . . . .	7
VII. References . . . . .	7
Appendix A . . . . .	8
Appendix B . . . . .	9

## FIGURES

		<u>Page</u>
Figure 1.	Time Interval of Periods Utilized in This Study . . . . .	10
Figure 2a-e.	Probability of Precipitation Events at PHX and FLG by Periods for Types 1-5 from 0000 GMT and 1200 GMT RAOBS . . . . .	11
Figure 3a-b.	Percent of Total Precipitation Events by Periods which Occurred in Types 1-4 from 0000 GMT and 1200 GMT RAOBS . . . . .	12
Figure 4a-b.	Graphical Representation of the Proba- bility of Precipitation Events by Periods at PHX and FLG for Types 1-5. RAOBS at 0000 GMT And 1200 GMT . . . . .	13
Figure 5a-d.	Breakdown Probabilities of Measurable Precipitation Events for the Presence or Lack of Moisture above 700 mb by Periods at PHX and FLG for Types 1-4. RAOBS at 0000 GMT and 1200 GMT . . . . .	14
Figure 6a-d.	Comparisons of Test Data Brier Scores Derived from Original (POPA), and Breakdown (POPB) Probabilities of Measurable Precipitation to those from Final MOS Probabilities of the Computer Run Concurrent with RAOB Time . . . . .	15
Figure 7a-d.	Comparisons of Test Data Brier Scores Derived from Original (POPA), and Breakdown (POPB) Probabilities of Measurable Precipitation to those from Final MOS Probabilities of the Computer Run 12 Hours Previous to RAOB Time . . . . .	16

THE DEPTH OF THE MARINE LAYER AT SAN DIEGO AS RELATED TO  
SUBSEQUENT COOL SEASON PRECIPITATION EPISODES IN ARIZONA

Ira S. Brenner  
Weather Service Forecast Office  
Phoenix\*, Arizona

ABSTRACT. The relationship between the depth of the marine layer at San Diego, California, and potential precipitation episodes in Arizona during the cool season is studied. It is shown that a marine layer from the surface to at least the 700-mb level is generally necessary for consideration of a subsequent widespread precipitation episode in Arizona. The relationship of the height of this marine inversion to the current vertical motion field is also discussed.

I. INTRODUCTION

It has long been subjectively recognized by Arizona forecasters that a correlation exists during the cool season (October-April) between the depth of the marine layer at San Diego (MYF) and potential precipitation episodes in Arizona. This is logical since MYF would be directly upstream from Arizona in the southwesterly flow preceding an advancing upper level trough.

It is felt that the height of the top of the marine layer is proportionate to the intensity of the vertical-motion field being superimposed on the area as an upper trough approaches. All too frequently, a vertical-motion field sufficiently strong to produce precipitation west of the coastal mountains is insufficient for widespread precipitation in Arizona. The theory being tested is that the vertical motion field west of the coastal range must be strong enough to raise the top of the marine layer to at least the 700-mb level in order to consider a widespread precipitation episode in Arizona. Numerous articles have been written on the subject of the quantitative effects of Positive Vorticity Advection (PVA) and subsequent vertical-motion fields. Two of the more pertinent articles for Arizona include Brenner (1979) and Rosendal (1976). However, it must be realized that the magnitude of the PVA/vertical-motion field necessary to deepen the MYF marine inversion through at least the 700-mb level during a given time interval will vary since, among other things, it will be a function of the available initial moisture values. The objective of this study was to demonstrate that the depth of the marine layer at MYF can be used on a real-time basis as a means for indirectly, but nevertheless reliably, measuring the relative magnitude of the current vertical-motion field ahead of an approaching upper trough.

During 1977 and 1978, an investigation was conducted to try and determine a more precise relationship between the observed vertical moisture profile at MYF and subsequent precipitation (as well as nonprecipitation) episodes in Arizona. Plotted data from balloon releases (RAOBS) dating from March

\*Present affiliation: National Hurricane Center, Miami, Florida

1971 through December 1975, were graciously loaned by the San Diego Weather Service Office for use as the developmental data base. The period of study involved data for the months of October through April only. Therefore, a total of thirty-three months of RAOBS comprised the developmental sample. With RAOBS for both 0000 and 1200 GMT available for nearly every day, approximately two thousand cases completed the developmental data base.

## II. DEVELOPMENTAL DATA BASE STRATIFICATION

Moisture distributions were sampled in terms of the summation of the temperature-dew point spread inventoried every 50 mb through a predetermined column:

$$(1) \quad A = (T-Td)_x + (T-Td)_{x-50} + \dots + (T-Td)_y$$

where  $(T-Td)$  is the temperature-dew point spread at a given level,  $x = 1000$  mb,  $y = 850$  mb.

$$(2) \quad B = (T-Td)_x + (T-Td)_{x-50} + \dots + (T-Td)_z$$

where  $(T-Td)$  is the temperature-dew point spread at a given level,  $x = 1000$  mb,  $z = 700$  mb.

The initial column stratifications (referred to as Types in the text) were selected as follows:

Type 1 equal  $A \leq 10^\circ\text{C}$  and  $B \leq 25^\circ\text{C}$ .

Type 2 equal  $A \leq 10^\circ\text{C}$  and  $B > 25^\circ\text{C}$ .

Type 3 equal  $11^\circ\text{C} \leq A \leq 30^\circ\text{C}$  and  $11^\circ\text{C} \leq B \leq 60^\circ$ .

Type 4 equal  $11^\circ\text{C} \leq A \leq 30^\circ\text{C}$  and  $B > 60^\circ\text{C}$ .

Type 5 All remaining cases.

Types 1 and 2 sampled all the available cases where a nearly saturated column of air existed in at least the lower 5000 feet. This was most frequently associated with a deep marine layer. Type 1 was designed to examine cases where the vertical-motion field was sufficiently strong to bring this layer through at least the 700-mb level, while Type 2 assumed the marine layer top short of 700 mb, but above 850 mb.

The theory being tested by Types 1 and 2, as mentioned earlier, was that the vertical-motion field west of the coastal range must be strong enough to deepen the marine layer through at least the 700-mb level in order to consider a widespread precipitation episode in Arizona. This same basic theory was tested by Types 3 and 4 also, but the required amounts of available moisture at MYF were scaled down. All remaining cases were included in Type 5.

### III. DEVELOPMENTAL DATA SAMPLE ANALYSIS

All the available MYF RAOBS from March 1971 through December 1975 (October through April only) were examined and separated into the various Types described earlier. Data from 0000 GMT RAOBS were analyzed apart from that of 1200 GMT. Data sheets for each Type were prepared and the dates of the respective RAOBS corresponding to each Type were recorded. Then four consecutive 12-hour periods (Figure 1) were individually examined for each date to determine if precipitation occurred at Phoenix (PHX, elevation 1100 ft) or Flagstaff (FLG, elevation 7000 ft). Throughout the remainder of this article, these four periods will simply be referred to as "Period 1, Period 2, etc.". However, when reference is made to the periods used by the National Weather Service for forecasts and comparison with Model Output Statistic (MOS) probabilities, the terms "FP Period 1, FP Period 2, etc." will be used. After tabulation, the number of measurable cases of precipitation only, and then the number of measurable and trace cases were totaled for each Type. Percent occurrences (in effect, conditional climatological probabilities for Periods 1-4 of this sample) were then computed for each. The results are shown in Figures 2a-e. Enough curiosity was raised to try and determine the percent of total measurable, as well as total measurable and trace cases that were caught by the combined Types 1-4. It was hoped that the percentage would be high enough to consider any adverse effects from rapidly changing conditions at balloon release times to be only an occasional compromising factor to the overall study. Figure 3 shows the results for Periods 1-4.

An analysis of Figure 3a reveals that in general, Types 1-4 for the 1200 GMT RAOBS caught on the order of 80-90% of the total measurable cases in the study for Period 1, 70-80% of the cases for Period 2, 60-70% for Period 3, and 50-60% for Period 4. The 0000 GMT RAOBS did not perform as well, indicating basically 70-80% for Period 1, 60-70% for Period 2, 50-60% for Period 3, and 40-50% for Period 4. An overall decrease in reliability occurred, as observed in Figure 3b, when the measurable and trace cases were considered. This was expected, since trace cases can frequently occur with middle and/or high-level moisture only. Perhaps another reason would be due to troughs approaching from a more northerly trajectory. Nevertheless, considering the overall rarity of precipitation events in Arizona and the fact that only one parameter (moisture at a fixed location) was being tested, it was felt that Types 1-4 locked in on those measurable events that did occur quite well. This was particularly true in Periods 1 and 2. One should be reminded at this point that this is only a climatological study, and although it would appear that this study has considerable prognostic value, it should be primarily viewed from a diagnostic standpoint when used operationally.

As mentioned earlier, 1200 GMT RAOBS outperformed the 0000 GMT RAOB data in the analysis of Figure 3. This diurnal conflict is intriguing. The 0000 GMT RAOBS had nearly 100 less cases per period in the total sample size for the combined Types 1-4 than the 1200 GMT RAOBS. Considerably more precipitation events occurred in Type 5 using 0000 GMT RAOBS as opposed to 1200 GMT data. One could speculate here that the problem is likely related to afternoon heating and mixing resulting in larger temperature-dew point spreads in the 1000- to 850-mb layer. Therefore, even though on a given day the vertical-motion field might still be strong enough

to give widespread precipitation in Arizona, the 0000 GMT RAOB may occasionally fail to satisfy the criteria for any of Types 1-4.

A graphical representation of the data presented in Figures 2a-e is shown in Figures 4a,b. Note that of the five Types for both PHX and FLG at 0000 GMT as well as 1200 GMT, the two Types involving high moisture values concentrated in at least the 1000-mb - 700-mb layer yielded the highest probabilities (Types 1 and 3). Types 2 and 4, which have high moisture only up to 850 mb, yielded lower probabilities (significantly lower for PHX) than those obtained by Types 1 and 3. This strongly suggests that high moisture values below 850 mbs, complimented by moisture in the 850-mb - 700-mb layer, is necessary for consideration of widespread precipitation in Arizona. This point is additionally supported by a comparison of Types 3 and 4. Type 4 involved the scaled-down moisture criteria in the 1000-mb - 850-mb layer and "dry" conditions between 850 mb and 700 mb. This, in itself, resulted in relatively low probabilities. The addition of moisture to the 850-mb - 700-mb layer to this, as shown in Type 4, with no change below 850 mb, sharply increased the probabilities (see the graphs of Type 3 for FLG and PHX).

Interest was then aroused as to the potential additional effects of high moisture above the 700-mb level. Therefore, Types 1 and 2 were tested for the effects of varying moisture supply above 700 mb. The criteria used was as follows:

$$(3) \quad C = (T-Td)_z + (T-Td)_{z-50} + \dots + (T-Td)_v$$

where (T-Td) is the temperature-dew point spread at a given level,  $z = 700$  mb,  $v = 400$  mb.

These were segregated such that:

- Sub-Type 1a equal Type 1 and  $C \leq 60^\circ\text{C}$ .
- Sub-Type 1b equal Type 1 and  $C > 60^\circ\text{C}$ .
- Sub-Type 2a equal Type 2 and  $C \leq 60^\circ\text{C}$ .
- Sub-Type 2b equal Type 2 and  $C > 60^\circ\text{C}$ .
- Sub-Type 3a equal Type 3 and  $C \leq 60^\circ\text{C}$ .
- Sub-Type 3b equal Type 3 and  $C > 60^\circ\text{C}$ .
- Sub-Type 4a equal Type 4 and  $C \leq 60^\circ\text{C}$ .
- Sub-Type 4b equal Type 4 and  $C > 60^\circ\text{C}$ .

Climatological probabilities for these Sub-Types (hereafter called "breakdown pops") were then derived and are displayed in Figure 5a-d with the original combined (or non-substratified) probabilities from Figure 2 a-e for comparison. The Sub-Typing resulted in data samples generally too small to be considered representative. Despite this, in most instances, the presence of high moisture values above 700 mb increased the probabilities from that of the original combined values, while the absence of this moisture had the opposite effect.



#### IV. INDEPENDENT TEST DATA

The months of October 1977 through April 1978 were utilized as test data. This provided a total sample size of 414 cases. Brier scores were totaled using the original combined pops for measurable precipitation (POPA) from Figures 2a-e as well as the breakdown pops (POPB). Comparisons were then made to the corresponding Final Model Output Statistics (MOS) Brier score in each of the three National Weather Service's FP periods. The results are listed in Figure 6a-d. Brier scores are rounded off and the decimal points displaced for convenience. As can be seen, the usage of the breakdown pops (POPB) generally degraded the results (increased the Brier scores) from those obtained by POPA. This was quite likely due to the problem of small sample size alluded to earlier. The breakdown pops did have a positive influence in a few cases. In general, the number of cases used to derive the breakdown probability for the presence of upper level moisture was too small to seriously consider the results reliable.

The comparison of POPA to MOS Brier scores displayed a few significant areas where the MOS forecasts could possibly be improved upon on an operational basis. Those listed below include periods where POPA Brier scores were less than or equal to the MOS Brier score in any period, or where POPA was less than 30 units above the MOS score in Periods 1 or 2 (indicating MOS was only slightly better than conditional climatology in the short term).

#### MOS WEAKNESS LIST #1

	RAOB TIME	STATION	TYPE	FP PERIOD	SAMPLE SIZE PER PERIOD
1.	0000GMT	PHX	2	1,2,3	11
2.	0000GMT	FLG	2	2,3	11
3.	0000GMT	PHX	1	2,3	12
4.	0000GMT	FLG	1	1,2,3	12
5.	1200GMT	PHX	1	1,2	20
6.	1200GMT	FLG	1	1,2	20
7.	0000GMT	PHX	4	1	26
8.	1200GMT	PHX	4	1,2,3	27
9.	1200GMT	PHX	3	1,2	11
10.	1200GMT	FLG	3	1	11

Of special interest here is that for Type 1, the wettest and most important of the Types in terms of precipitation events, MOS commonly was only slightly better or actually worse than the conditional climatological pops from the study (POPA). This was true at both RAOB times and for both PHX and FLG. A review of the appropriate data indicated that MOS had a definite tendency for forecasting rather low probabilities (0-30%)--many on which precipitation occurred.

A more detailed examination of the possible weaknesses in the MOS forecasts for the above MOS WEAKNESS LIST #1 is found in Appendix A and labeled WEAKNESS LIST #1. The numbers of the 1-10 in WEAKNESS LIST #1 correspond to the same numbers in the above MOS WEAKNESS LIST #1.

A return to Figure 3 brings forth another interesting point. The best results for measurable as well as measurable plus trace cases were in Periods 1 and 2. Perhaps the operational forecaster could also be served by this study in terms of an updating tool. The analysis of Brier scores just examined from Figure 6 involved a comparison of data from a given RAOB to the MOS run from the same time as the RAOB. However, the RAOB is nearly 10 hours old by the time the first FP period begins. In actuality, a given RAOB is received almost at the beginning time of the first FP period MOS probabilities from the previous run. For example, the 1200GMT RAOB is received and plotted by the time the first FP period MOS pop from the previous 0000GMT run is only about 2 hours old. The utility of this study, examined from the standpoint of an updating aid, is tabulated in Figure 7a-d.

The comparison of POPA to MOS Brier scores for purposes of updating also indicated areas where MOS forecasts from the previous runs were potentially weak. Opportunities for improvement upon MOS forecasts existed in the following categories:

MOS WEAKNESS LIST #2

	<u>RAOB TIME</u>	<u>STATION</u>	<u>TYPE</u>	<u>FP PERIOD</u>	<u>SAMPLE SIZE PER PERIOD</u>
1.	0000GMT	PHX	2	1,2,3	11
2.	0000GMT	FLG	2	1,2,3	11
3.	0000GMT	PHX	1	1,2,3	12
4.	0000GMT	FLG	1	1,2,3	12
5.	1200GMT	PHX	1	1,2	20
6.	1200GMT	FLG	1	1,2,3	20
7.	0000GMT	PHX	4	2,3	26
8.	0000GMT	FLG	4	1, 3	26
9.	1200GMT	PHX	4	1,2,3	27
10.	1200GMT	FLG	4	1	27
11.	1200GMT	PHX	3	2,3	11
12.	1200GMT	FLG	3	3	11

Note here also that Type 1 showed up again at both RAOB times and for both PHX and FLG. As with MOS WEAKNESS LIST #1, an investigation of the data revealed that the MOS tendency to forecast rather low probabilities, on which precipitation occurred, persisted. Appendix B uses a format similar to Appendix A for further describing the possible MOS weaknesses corresponding to MOS WEAKNESS LIST #2.

## V. CONCLUSIONS

This conditional climatological study was considered to be a beneficial diagnostic forecast aid for PHX WSFO. Even if used strictly from an objective standpoint, the study yielded excellent results. The added usage of a limited amount of subjective reasoning and modification will improve the operational results even further. The utility of the study extended beyond the capacity of making three-period probability forecasts. It was found that the study also served the forecaster successfully as an updating tool.

The developmental data sample did not stratify precipitation episodes at PHX or FLG by storm origin or trajectory. Despite this, results, particularly using 1200GMT MYF RAOBS, still displayed a definite relationship between the depth of the marine layer at MYF and subsequent widespread precipitation episodes in Arizona. It is felt that on an operational basis, subjective evaluation can be made to the study probabilities for cases where storms approach from a more northerly direction or when an unusually strong influx of tropical moisture is involved.

This investigation gave strong supportive evidence that high moisture content (i.e., a high marine inversion) at MYF from the surface to at least 700 mb is generally necessary for widespread precipitation episodes in Arizona. It is the opinion of the author that when the top of the marine layer at MYF is lifted to at least the 700-mb level ahead of an upper-level trough, the vertical-motion field will generally remain strong enough to produce widespread precipitation upon reaching Arizona.

Although not conclusive from this study, it would appear subjectively that additional high moisture values in the 700-mb - 400-mb layer enhance the probabilities of precipitation even further.

A fringe benefit of this study was the identification of potentially weak areas in the MOS probability forecasts. With reasonable discretion, forecasters can successfully use the results listed in the INDEPENDENT TEST DATA section to identify and hopefully improve upon available corresponding MOS forecasts.

## VI. ACKNOWLEDGMENTS

Appreciation is extended to the San Diego Weather Service Office for use of their plotted RAOBS and to Mrs. Tommie McCabe and Mrs. Evelyn Allan for their conscientious typing efforts.

## VII. REFERENCES

- Brenner, I. S., 1979: A Case Study of Rapid Air Mass Modification Over Arizona Due to Large-Scale Upward Motion. National Weather Digest, Vol. 4, No. 1, February, pp 18-26.
- Rosendal, Hans E., 1976: The Zero Relative Vorticity Line at 500 mb as Related to Precipitation and Surface Fronts. Manuscript from AMS Sixth Conference on Weather Forecasting and Analysis, Albany, New York.

APPENDIX A

WEAKNESS LIST #1

An analysis of the data from MOS WEAKNESS LIST #1 revealed the following information (Numbers 1-10 refer to the corresponding numbers in MOS WEAKNESS LIST #1):

1. MOS frequently forecast pops of 30% or greater with no cases of measurable precipitation occurring.
2. Measurable precipitation fell on more than half the cases when MOS forecast pops between and including 5% and 20%. Also, no measurable precipitation fell on MOS pops of 70% or greater.
3. No measurable precipitation fell on MOS pops of 70% or greater. Also measurable precipitation fell on half the cases where MOS pops were 20% or less.
4. A large amount of measurable precipitation events occurred on MOS pops of 30% or less.
5. Measurable precipitation fell on one half of the cases where MOS forecast pops of 5% - 20%.
6. Measurable precipitation fell on most of the cases where MOS forecast pops of 5% - 30%.
7. No measurable precipitation fell on MOS pops of 30% or greater.
8. No measurable precipitation fell on MOS pops of 20% - 60%.
9. Measurable precipitation fell on a 0% in FP Period 1. No measurable precipitation fell on MOS pops of 50% or greater in FP Period 2.
10. Several incidents where measurable precipitation fell were on MOS pops of 20% or less.

## APPENDIX B

An analysis of this data from MOS WEAKNESS LIST #2 revealed the following:

1. MOS frequently forecast pops of 30% or greater, and, excepting one case of .08 in. on a 60% pop, no other measurable precipitation occurred. For FP Period 1, MOS got precipitation on half of the 0% and 5%.
2. Measurable precipitation fell on more than half the cases when MOS forecast pops from 5% and 20%. Also, no measurable precipitation fell on MOS pops 70% or greater in FP Period 3.
3. No measurable precipitation fell 2 out of 3 times on MOS pops of 70% or greater. Measurable precipitation fell on half of the pops of 20% or less.
4. A large amount of measurable precipitation events occurred on MOS pops of 30% or less.
5. Measurable precipitation fell on half the cases where MOS forecast pops of 5% to 20%.
6. Measurable precipitation fell on most cases where MOS forecast pops of 5% to 30%.
7. No measurable precipitation fell on MOS pops of 40% or greater.
8. No measurable precipitation fell on MOS pops 50% or greater. Measurable precipitation fell on half the cases of MOS pops of 2%.
9. No measurable precipitation fell, excepting one case of .01 inch on a 20% MOS pop, on MOS pops in the 20-60% bracket.
10. No measurable precipitation fell, excepting one case, on MOS pops 50% or greater.
11. Measurable precipitation fell on a 0% in FP Period 2. No measurable precipitation fell on MOS pops of 50% or greater in FP Period 3.
12. No measurable precipitation fell on MOS pops of 50% or greater.

0000 GMT RAOB		
PERIOD NUMBER USED IN STUDY	TIME INTERVAL (GMT)	CORRESPONDING FP PERIOD
1	0000 - 1200	
2	1200 - 0000	1
3	0000 - 1200	2
4	1200 - 0000	3

1200 GMT RAOB		
PERIOD NUMBER USED IN STUDY	TIME INTERVAL (GMT)	CORRESPONDING FP PERIOD
1	1200 - 0000	
2	0000 - 1200	1
3	1200 - 0000	2
4	0000 - 1200	3

Figure 1. Time Interval of Periods Utilized in this Study.

0000 GMT RAOB

TYPE 1	STN PD	PHX				SAMPLE SIZE FOR TYPE 1	FLG			
		1	2	3	4		1	2	3	4
MEASURABLE CASES		11	9	9	4	25	18	19	17	12
MEASURABLE AND TRACE CASES		15	12	10	5	25	23	21	20	13
PERCENT MEASURABLE		44	36	36	16		72	76	68	48
PERCENT MEASURABLE AND TRACE		60	48	40	20		92	84	80	52

0000 GMT RAOB

TYPE 2	STN PD	PHX				SAMPLE SIZE FOR TYPE 2	FLG			
		1	2	3	4		1	2	3	4
MEASURABLE CASES		4	9	5	5	37	22	22	20	12
MEASURABLE AND TRACE CASES		14	16	11	8	37	27	25	23	15
PERCENT MEASURABLE		11	24	14	14		59	59	54	32
PERCENT MEASURABLE AND TRACE		38	43	30	22		73	68	62	41

0000 GMT RAOB

TYPE 3	STN PD	PHX				SAMPLE SIZE FOR TYPE 3	FLG			
		1	2	3	4		1	2	3	4
MEASURABLE CASES		20	11	8	8	50	35	30	20	22
MEASURABLE AND TRACE CASES		27	17	14	12	50	40	36	27	27
PERCENT MEASURABLE		40	22	16	16		70	60	40	44
PERCENT MEASURABLE AND TRACE		54	34	28	24		80	72	54	54

1200 GMT RAOB

TYPE 1	STN PD	PHX				SAMPLE SIZE FOR TYPE 1	FLG			
		1	2	3	4		1	2	3	4
MEASURABLE CASES		19	16	6	6	41	35	27	19	15
MEASURABLE AND TRACE CASES		25	22	11	8	41	40	33	23	23
PERCENT MEASURABLE		46	39	15	15		85	66	46	37
PERCENT MEASURABLE AND TRACE		61	54	27	20		98	81	56	56

1200 GMT RAOB

TYPE 2	STN PD	PHX				SAMPLE SIZE FOR TYPE 2	FLG			
		1	2	3	4		1	2	3	4
MEASURABLE CASES		13	17	17	15	112	55	46	47	40
MEASURABLE AND TRACE CASES		25	28	30	22	112	74	64	57	50
PERCENT MEASURABLE		12	15	15	13		49	41	42	36
PERCENT MEASURABLE AND TRACE		22	25	27	20		66	57	51	45

1200 GMT RAOB

TYPE 3	STN PD	PHX				SAMPLE SIZE FOR TYPE 3	FLG			
		1	2	3	4		1	2	3	4
MEASURABLE CASES		13	8	9	7	42	33	26	14	11
MEASURABLE AND TRACE CASES		26	19	13	11	42	34	35	21	15
PERCENT MEASURABLE		31	19	21	17		79	62	33	26
PERCENT MEASURABLE AND TRACE		62	45	31	26		81	83	50	36

0000 GMT RAOB

TYPE 4	STN PD	PHX				SAMPLE SIZE FOR TYPE 4	FLG			
		1	2	3	4		1	2	3	4
MEASURABLE CASES		16	12	14	9	109	35	37	33	31
MEASURABLE AND TRACE CASES		23	23	28	18	109	51	52	44	37
PERCENT MEASURABLE		15	11	13	8		32	34	30	28
PERCENT MEASURABLE AND TRACE		21	21	26	17		47	48	40	34

0000 GMT RAOB

TYPE 5	STN PD	PHX				SAMPLE SIZE FOR TYPE 5	FLG			
		1	2	3	4		1	2	3	4
MEASURABLE CASES		14	20	27	38	781	50	63	73	90
MEASURABLE AND TRACE CASES		41	42	52	66	781	93	98	125	133
PERCENT MEASURABLE		2	3	4	5		6	8	9	12
PERCENT MEASURABLE AND TRACE		5	5	7	8		12	13	16	17

1200 GMT RAOB

TYPE 4	STN PD	PHX				SAMPLE SIZE FOR TYPE 4	FLG			
		1	2	3	4		1	2	3	4
MEASURABLE CASES		7	4	7	6	123	23	23	29	30
MEASURABLE AND TRACE CASES		11	17	16	17	123	34	36	37	41
PERCENT MEASURABLE		6	3	6	5		19	19	24	24
PERCENT MEASURABLE AND TRACE		9	14	13	14		28	29	30	33

1200 GMT RAOB

TYPE 5	STN PD	PHX				SAMPLE SIZE FOR TYPE 5	FLG			
		1	2	3	4		1	2	3	4
MEASURABLE CASES		7	15	24	25	684	28	38	63	63
MEASURABLE AND TRACE CASES		23	29	44	51	684	50	73	92	104
PERCENT MEASURABLE		1	2	4	4		4	6	9	9
PERCENT MEASURABLE AND TRACE		3	4	6	8		7	11	13	15

Figures 2a-e: Probability of Precipitation Events at PHX and FLG by Periods for Types 1-5 from 0000 GMT and 1200 GMT RAOBS.

RELIABILITY  
MEASURABLE PRECIPITATION  
FIGURE 3A

	PERIOD	1200 GMT				TOTAL SAMPLE SIZE PER PD	PERIOD	0000 GMT				TOTAL SAMPLE SIZE PER PD
		1	2	3	4			1	2	3	4	
TOTAL NUMBER OF EVENTS CAUGHT BY TYPES 1-4	PHX	52	45	39	34	318	PHX	51	41	36	26	221
	FLG	146	122	109	96		FLG	110	109	90	77	
TOTAL NUMBER OF EVENTS IN STUDY	PHX	59	60	63	59	1002	PHX	65	61	63	64	1002
	FLG	174	160	172	159		FLG	160	172	163	167	
PERCENT OF TOTAL EVENTS WHICH OCCURRED IN TYPES 1-4	PHX	88%	75%	62%	58%		PHX	78%	67%	57%	41%	
	FLG	84%	76%	63%	60%		FLG	69%	63%	55%	46%	

RELIABILITY  
MEASURABLE AND TRACE  
FIGURE 3B

	PERIOD	1200 GMT				TOTAL SAMPLE SIZE PER PD	PERIOD	0000 GMT				TOTAL SAMPLE SIZE PER PD
		1	2	3	4			1	2	3	4	
TOTAL NUMBER OF EVENTS CAUGHT BY TYPES 1-4	PHX	87	86	70	58	318	PHX	79	68	63	43	221
	FLG	182	168	138	129		FLG	141	134	114	92	
TOTAL NUMBER OF EVENTS IN STUDY	PHX	110	115	114	109	1002	PHX	120	110	115	109	1002
	FLG	232	241	230	235		FLG	234	232	239	225	
PERCENT OF TOTAL EVENTS WHICH OCCURRED IN TYPES 1-4	PHX	79%	75%	61%	53%		PHX	66%	62%	55%	39%	
	FLG	78%	70%	60%	55%		FLG	60%	58%	48%	41%	

FIGURES 3A and 3B: Probability of Precipitation Events at PHX and FLG by Periods for Types 1-5 from 0000 GMT and 1200 GMT RAOBS.







Figure 4a-b: Graphical Representation of the Probability of Precipitation Events by Periods at PHX and FLG for Types 1-5. RAOBS at 0000 GMT and 1200 GMT.

PROBABILITY OF MEASURABLE PRECIPITATION

TYPE 1

PHX

0000 GMT						1200 GMT					
	N	PD1	PD2	PD3	PD4		N	PD1	PD2	PD3	PD4
WITH UPPER LEVEL MOISTURE	14	50	50	36	21	WITH UPPER LEVEL MOISTURE	13	62	54	23	15
COMBINED	25	44	36	36	16	COMBINED	41	46	39	15	15
WITHOUT UPPER LEVEL MOISTURE	11	36	18	36	9	WITHOUT UPPER LEVEL MOISTURE	28	39	32	11	14

FLG											
0000 GMT						1200 GMT					
	N	PD1	PD2	PD3	PD4		N	PD1	PD2	PD3	PD4
WITH UPPER LEVEL MOISTURE	14	79	79	64	50	WITH UPPER LEVEL MOISTURE	13	92	69	54	38
COMBINED	25	72	76	68	48	COMBINED	41	85	66	46	37
WITHOUT UPPER LEVEL MOISTURE	11	64	73	73	45	WITHOUT UPPER LEVEL MOISTURE	28	82	64	43	36

PROBABILITY OF MEASURABLE PRECIPITATION

TYPE 2

PHX

0000 GMT						1200 GMT					
	N	PD1	PD2	PD3	PD4		N	PD1	PD2	PD3	PD4
WITH UPPER LEVEL MOISTURE	7	43	71	14	43	WITH UPPER LEVEL MOISTURE	10	20	40	20	0
COMBINED	37	11	24	14	14	COMBINED	112	12	15	15	13
WITHOUT UPPER LEVEL MOISTURE	30	3	20	13	7	WITHOUT UPPER LEVEL MOISTURE	102	12	13	15	15

FLG											
0000 GMT						1200 GMT					
	N	PD1	PD2	PD3	PD4		N	PD1	PD2	PD3	PD4
WITH UPPER LEVEL MOISTURE	7	86	86	71	57	WITH UPPER LEVEL MOISTURE	10	60	60	60	40
COMBINED	37	59	59	54	32	COMBINED	112	49	41	42	36
WITHOUT UPPER LEVEL MOISTURE	30	53	53	50	27	WITHOUT UPPER LEVEL MOISTURE	102	48	39	40	35

PROBABILITY OF MEASURABLE PRECIPITATION

TYPE 3

PHX

0000 GMT						1200 GMT					
	N	PD1	PD2	PD3	PD4		N	PD1	PD2	PD3	PD4
WITH UPPER LEVEL MOISTURE	10	40	20	20	10	WITH UPPER LEVEL MOISTURE	15	27	27	47	27
COMBINED	50	40	22	16	16	COMBINED	42	31	19	21	17
WITHOUT UPPER LEVEL MOISTURE	40	40	23	15	18	WITHOUT UPPER LEVEL MOISTURE	27	33	15	7	11

FLG											
0000 GMT						1200 GMT					
	N	PD1	PD2	PD3	PD4		N	PD1	PD2	PD3	PD4
WITH UPPER LEVEL MOISTURE	10	90	80	30	30	WITH UPPER LEVEL MOISTURE	15	87	73	47	40
COMBINED	50	70	60	40	44	COMBINED	42	79	62	33	26
WITHOUT UPPER LEVEL MOISTURE	40	65	55	43	48	WITHOUT UPPER LEVEL MOISTURE	27	74	56	26	19

PROBABILITY OF MEASURABLE PRECIPITATION

TYPE 4

PHX

0000 GMT						1200 GMT					
	N	PD1	PD2	PD3	PD4		N	PD1	PD2	PD3	PD4
WITH UPPER LEVEL MOISTURE	9	33	56	33	11	WITH UPPER LEVEL MOISTURE	3	33	33	33	0
COMBINED	109	15	11	13	8	COMBINED	123	6	3	6	5
WITHOUT UPPER LEVEL MOISTURE	100	13	7	11	8	WITHOUT UPPER LEVEL MOISTURE	120	5	3	5	5

FLG											
0000 GMT						1200 GMT					
	N	PD1	PD2	PD3	PD4		N	PD1	PD2	PD3	PD4
WITH UPPER LEVEL MOISTURE	9	56	67	44	44	WITH UPPER LEVEL MOISTURE	3	33	67	67	67
COMBINED	109	32	34	30	28	COMBINED	123	19	19	24	24
WITHOUT UPPER LEVEL MOISTURE	100	30	31	29	27	WITHOUT UPPER LEVEL MOISTURE	120	18	18	23	23

FIGURE 5A-D: BREAKDOWN PROBABILITIES OF MEASURABLE PRECIPITATION EVENTS FOR THE PRESENCE OR LACK OF MOISTURE ABOVE 700 MB BY PERIODS AT PHX AND FLG FOR TYPES 1-4. RAOBS AT 0000 GMT AND 1200 GMT.

COMPARISON OF TOTAL BRIER SCORES FOR BOTH ORIGINAL POPS  
(POPA) AND BREAKDOWN POPS (POPB) TO CURRENT MOS POPS

CURRENT 0000 GMT RAOB  
CURRENT 0000 GMT MOS

TYPE 1 - N=12

PHX			FLG		
PD1 (FP)	PD2 (FP)	PD3 (FP)	PD1 (FP)	PD2 (FP)	PD3 (FP)
POPA	POPA	POPA	POPA	POPA	POPA
372   MOS	332   MOS	288   MOS	168   MOS	228   MOS	300   MOS
POPB   209	POPB   345	POPB   324	POPB   217	POPB   299	POPB   388
495	332	313	193	237	300

CURRENT 0000 GMT RAOB  
CURRENT 0000 GMT MOS

TYPE 2 - N=11

PHX			FLG		
PD1 (FP)	PD2 (FP)	PD3 (FP)	PD1 (FP)	PD2 (FP)	PD3 (FP)
POPA	POPA	POPA	POPA	POPA	POPA
104   MOS	91   MOS	91   MOS	276   MOS	275   MOS	259   MOS
POPB   100	POPB   125	POPB   182	POPB   238	POPB   396	POPB   301
104	91	91	276	275	259

COMPARISON OF TOTAL BRIER SCORES FOR BOTH ORIGINAL POPS  
(POPA) AND BREAKDOWN POPS (POPB) TO CURRENT MOS POPS

CURRENT 0000 GMT RAOB  
CURRENT 0000 GMT MOS

TYPE 3 - N=25

PHX			FLG		
PD1 (FP)	PD2 (FP)	PD3 (FP)	PD1 (FP)	PD2 (FP)	PD3 (FP)
POPA	POPA	POPA	POPA	POPA	POPA
520   MOS	520   MOS	400   MOS	600   MOS	600   MOS	540   MOS
POPB   294	POPB   346	POPB   231	POPB   356	POPB   301	POPB   230
580	520	433	672	617	681

CURRENT 0000 GMT RAOB  
CURRENT 0000 GMT MOS

TYPE 4 - N=26

PHX			FLG		
PD1 (FP)	PD2 (FP)	PD3 (FP)	PD1 (FP)	PD2 (FP)	PD3 (FP)
POPA	POPA	POPA	POPA	POPA	POPA
106   MOS	266   MOS	266   MOS	354   MOS	394   MOS	474   MOS
POPB   121	POPB   176	POPB   124	POPB   185	POPB   227	POPB   223
58	242	266	354	388	468

-15-

COMPARISON OF TOTAL BRIER SCORES FOR BOTH ORIGINAL POPS  
(POPA) AND BREAKDOWN POPS (POPB) TO CURRENT MOS POPS

CURRENT 1200 GMT RAOB  
CURRENT 1200 GMT MOS

TYPE 1 - N=20

PHX			FLG		
PD1 (FP)	PD2 (FP)	PD3 (FP)	PD1 (FP)	PD2 (FP)	PD3 (FP)
POPA	POPA	POPA	POPA	POPA	POPA
520   MOS	500   MOS	380   MOS	340   MOS	500   MOS	500   MOS
POPB   567	POPB   483	POPB   366	POPB   434	POPB   476	POPB   352
500	510	450	350	530	500

CURRENT 1200 GMT RAOB  
CURRENT 1200 GMT MOS

TYPE 2 - N=17

PHX			FLG		
PD1 (FP)	PD2 (FP)	PD3 (FP)	PD1 (FP)	PD2 (FP)	PD3 (FP)
POPA	POPA	POPA	POPA	POPA	POPA
368   MOS	428   MOS	257   MOS	412   MOS	392   MOS	392   MOS
POPB   251	POPB   175	POPB   236	POPB   179	POPB   147	POPB   226
372	428	284	412	372	372

COMPARISON OF TOTAL BRIER SCORES FOR BOTH ORIGINAL POPS  
(POPA) AND BREAKDOWN POPS (POPB) TO CURRENT MOS POPS

CURRENT 1200 GMT RAOB  
CURRENT 1200 GMT MOS

TYPE 3 - N=11

PHX			FLG		
PD1 (FP)	PD2 (FP)	PD3 (FP)	PD1 (FP)	PD2 (FP)	PD3 (FP)
POPA	POPA	POPA	POPA	POPA	POPA
164   MOS	164   MOS	224   MOS	316   MOS	179   MOS	299   MOS
POPB   147	POPB   172	POPB   191	POPB   291	POPB   118	POPB   241
169	207	219	329	195	316

CURRENT 1200 GMT RAOB  
CURRENT 1200 GMT MOS

TYPE 4 - N=27

PHX			FLG		
PD1 (FP)	PD2 (FP)	PD3 (FP)	PD1 (FP)	PD2 (FP)	PD3 (FP)
POPA	POPA	POPA	POPA	POPA	POPA
97   MOS	97   MOS	97   MOS	408   MOS	408   MOS	288   MOS
POPB   127	POPB   137	POPB   199	POPB   352	POPB   257	POPB   279
97	97	97	408	408	288

FIGURE 6a-d. COMPARISONS OF TEST DATA BRIER SCORES DERIVED FROM ORIGINAL (POPA) AND BREAKDOWN (POPB) PROBABILITIES OF MEASURABLE PRECIPITATION TO THOSE FROM FINAL MOS PROBABILITIES OF THE COMPUTER RUN CONCURRENT WITH RAOB TIME.

COMPARISON OF TOTAL BRIER SCORES FOR BOTH ORIGINAL POPS  
(POPA) AND BREAKDOWN POPS (POPB) TO PREVIOUS MOS POPS

CURRENT 0000 GMT RAOB  
CURRENT 0000 GMT MOS

TYPE 1 - N=12

PHX			FLG		
PD1	PD2(FP PD1)	PD3(FP PD2)	PD1	PD2(FP PD1)	PD3(FP PD2)
POPA	POPA	POPA	POPA	POPA	POPA
372	372	332	148	168	228
MOS	MOS	MOS	MOS	MOS	MOS
POPB	POPB	POPB	POPB	POPB	POPB
590	374	332	509	301	289
335	495	332	128	193	237

CURRENT 0000 GMT RAOB  
CURRENT 1200 GMT MOS

TYPE 2 - N=11

PHX			FLG		
PD1	PD2(FP PD1)	PD3(FP PD2)	PD1	PD2(FP PD1)	PD3(FP PD2)
POPA	POPA	POPA	POPA	POPA	POPA
251	104	91	236	276	275
MOS	MOS	MOS	MOS	MOS	MOS
POPB	POPB	POPB	POPB	POPB	POPB
222	142	158	518	318	324
288	104	91	275	276	275

COMPARISON OF TOTAL BRIER SCORES FOR BOTH ORIGINAL POPS  
(POPA) AND BREAKDOWN POPS (POPB) TO PREVIOUS MOS POPS

CURRENT 0000 GMT RAOB  
PREVIOUS 1200 GMT MOS

TYPE 3 - N=25

PHX			FLG		
PD1	PD2(FP PD1)	PD3(FP PD2)	PD1	PD2(FP PD1)	PD3(FP PD2)
POPA	POPA	POPA	POPA	POPA	POPA
520	520	520	785	600	600
MOS	MOS	MOS	MOS	MOS	MOS
POPB	POPB	POPB	POPB	POPB	POPB
436	411	446	376	447	431
520	580	520	913	672	617

CURRENT 0000 GMT RAOB  
PREVIOUS 1200 GMT MOS

TYPE 4 - N=26

PHX			FLG		
PD1	PD2(FP PD1)	PD3(FP PD2)	PD1	PD2(FP PD1)	PD3(FP PD2)
POPA	POPA	POPA	POPA	POPA	POPA
284	106	266	336	354	394
MOS	MOS	MOS	MOS	MOS	MOS
POPB	POPB	POPB	POPB	POPB	POPB
131	159	282	414	262	394
282	58	242	348	354	388

-16-

COMPARISON OF TOTAL BRIER SCORES FOR BOTH ORIGINAL POPS  
(POPA) AND BREAKDOWN POPS (POPB) TO PREVIOUS MOS POPS

CURRENT 1200 GMT RAOB  
PREVIOUS 0000 GMT MOS

TYPE 1 - N=20

PHX			FLG		
PD1	PD2(FP PD1)	PD3(FP PD2)	PD1	PD2(FP PD1)	PD3(FP PD2)
POPA	POPA	POPA	POPA	POPA	POPA
500	520	500	260	340	500
MOS	MOS	MOS	MOS	MOS	MOS
POPB	POPB	POPB	POPB	POPB	POPB
481	581	180	329	207	130
320	372	428	416	412	372

CURRENT 1200 GMT RAOB  
PREVIOUS 0000 GMT MOS

TYPE 2 - N=17

PHX			FLG		
PD1	PD2(FP PD1)	PD3(FP PD2)	PD1	PD2(FP PD1)	PD3(FP PD2)
POPA	POPA	POPA	POPA	POPA	POPA
337	368	428	425	412	392
MOS	MOS	MOS	MOS	MOS	MOS
POPB	POPB	POPB	POPB	POPB	POPB
180	259	180	329	207	130
320	372	428	416	412	372

COMPARISON OF TOTAL BRIER SCORES FOR BOTH ORIGINAL POPS  
(POPA) AND BREAKDOWN POPS (POPB) TO PREVIOUS MOS POPS

CURRENT 1200 GMT RAOB  
PREVIOUS 0000 GMT MOS

TYPE 3 - N=11

PHX			FLG		
PD1	PD2(FP PD1)	PD3(FP PD2)	PD1	PD2(FP PD1)	PD3(FP PD2)
POPA	POPA	POPA	POPA	POPA	POPA
219	164	164	284	316	179
MOS	MOS	MOS	MOS	MOS	MOS
POPB	POPB	POPB	POPB	POPB	POPB
106	146	231	163	223	181
219	169	207	291	329	195

CURRENT 1200 GMT RAOB  
PREVIOUS 0000 GMT MOS

TYPE 4 - N=17

PHX			FLG		
PD1	PD2(FP PD1)	PD3(FP PD2)	PD1	PD2(FP PD1)	PD3(FP PD2)
POPA	POPA	POPA	POPA	POPA	POPA
7	97	97	228	408	408
MOS	MOS	MOS	MOS	MOS	MOS
POPB	POPB	POPB	POPB	POPB	POPB
43	117	145	310	365	292
7	97	97	228	408	408

FIGURE 7a-d. COMPARISONS OF TEST DATA BRIER SCORES DERIVED FROM ORIGINAL (POPA) AND BREAKDOWN (POPB) PROBABILITIES OF MEASURABLE PRECIPITATION TO THOSE FROM FINAL MOS PROBABILITIES OF THE COMPUTER RUN 12 HOURS PREVIOUS TO RAOB TIME.

NOAA Technical Memoranda NWSNR: (Continued)

- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (60N-74-11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (60N-74-11407/AS)
- 94 Conditional Probability of Visibility Loss than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (60N-74-11555/AS)
- 96 Map Type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (60N-75-10428/AS)
- 97 Eastern Pacific Cut-off Low of April 21-23, 1974. William J. Alder and George R. Miller, January 1976. (PB-250-711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (60N-75-10719/AS)
- 99 A Study of Flash Flood Susceptibility--A Basin in Southern Arizona. Gerald Williams, August 1976. (60N-75-11368/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1976. (PB-246-902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB-253-853/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB-252-868/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruseha, Jr., February 1976. (PB-254-650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John G. Plankinton, Jr., March 1976. (PB-254-649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB-259-594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB-260-437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, Sept. 1976. (PB-273-877/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denny, November 1976. (PB-264-855/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB-265-941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California; Using Multiple Regression. Michael J. Card, February 1977. (PB-273-694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB-273-876/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB-268-847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-value. R. F. Quiring, April 1977. (PB-272-831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB-268-740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Barryl Randserson, June 1977. (PB-271-200/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R. F. Quiring, June 1977. (PB-271-704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB-271-742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB-272-661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB-275-155/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth E. Mielke, February 1978. (PB-281-567/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudge, April 1978. (PB-283-880/AS)
- 129 Fire Whirls. David W. Coans, May 1978. (PB-283-866/AS)
- 130 Flash-Flood Procedure. Ralph O. Hatch and Gerald Williams, May 1978. (PB-283-914/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olson, September 1978. (PB-289-916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R. G. Pappas, R. Y. Leo, and B. W. Fluke, October 1978. (PB-289-767/AS)
- 135 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB-291-317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB-291-195/AS)
- 133 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB-292-716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB-292-247/AS)
- 137 LFM 24-hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruseha, Jr., January 1979.
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979.
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979.
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather District. James Holcomb, April 1979.
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979.
- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979.

## NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

NOAA, the *National Oceanic and Atmospheric Administration*, was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth, and to assess the socioeconomic impact of natural and technological changes in the environment.

The six Major Line Components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

**PROFESSIONAL PAPERS**—Important definitive research results, major techniques, and special investigations.

**TECHNICAL REPORTS**—Journal quality with extensive details, mathematical developments, or data listings.

**TECHNICAL MEMORANDUMS**—Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

**CONTRACT AND GRANT REPORTS**—Reports prepared by contractors or grantees under NOAA sponsorship.

**TECHNICAL SERVICE PUBLICATIONS**—These are publications containing data, observations, instructions, etc. A partial listing: Data serials; Prediction and outlook periodicals; Technical manuals, training papers, planning reports, and information serials; and Miscellaneous technical publications.

**ATLAS**—Analysed data generally presented in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.



*Information on availability of NOAA publications can be obtained from:*

**ENVIRONMENTAL SCIENCE INFORMATION CENTER  
ENVIRONMENTAL DATA SERVICE  
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION  
U.S. DEPARTMENT OF COMMERCE**

**3300 Whitehaven Street, N.W.  
Washington, D.C. 20235**